Indicators	2024	2025	2026
KP-1d Percentage of people who inject drugs reached with HIV prevention programs - defined package of services	5200	5900	6600
HTS-3d Percentage of people who inject drugs that have received an HIV test during the reporting period in KP-specific programs and know their results	15502	16478	17482
KP-1c Percentage of sex workers reached with HIV prevention programs - defined package of services	1200	1500	1800
HTS-3c Percentage of sex workers that have received an HIV test during the reporting period in KP-specific programs and know their results	5393	5662	5946
KP-1a Percentage of men who have sex with men reached with HIV prevention programs - defined package of services	7300	8700	10000
HTS-3a Percentage of MSM that have received an HIV test during the reporting period in KP-specific programs and know their results	9312	9962	10724
KP-1b Percentage of transgender people reached with HIV prevention programs - defined package of services	5250	5800	6450
HTS-3b Percentage of TG that have received an HIV test during the reporting period in KP-specific programs and know their results	4119	4501	4966
KP-6a Number of MSM who received any PrEP product at least once during the reporting period	2599	2989	3436
KP-6d Number of PWID who received any PrEP product at least once during the reporting period	561	648	746
KP-6b Number of transgender people who received any PrEP product at least once during the reporting period	20	25	30
KP-6c Number of sex workers who received any PrEP product at least once during the reporting period	253	292	321
TBDT-1 Number of patients with of all forms of TB notified (i.e., bacteriologically confirmed + clinically diagnosed); *includes only those with new and relapse TB	600	156	
DRTB - Other 1 Number of RR-TB and/or MDR-TB cases covered with comprehensive psychosocial support	2374	1744	1744
DRTB-9 Treatment success rate of RR-TB and/or MDR-TB: Percentage of patients with RR and/or MDR-TB successfully treated	67%	69%	71%
DRTB-4 Percentage of people with RR-TB/MDR-TB who did not start treatment and/or started on treatment for MDR- TB who were lost to follow up during the first six months of treatment	6%	5.5%	5%
KP-5 Percentage of individuals receiving opioid substitution therapy who received treatment for at least 6 months	90%	91%	92%
TCS - Other 2: Percentage of HIV positive PWID linked to HIV care	95%	95%	95%